




**Neurotrophic Keratitis:  
Rare or Hiding in Plain  
Sight**  
Douglas K Devries, OD  
Eye Care Associates of Nevada

1

**Douglas K Devries, OD  
Disclosures**  
All Conflicts Have Been Mitigated

<ul style="list-style-type: none"> <li>Allergan Advisor</li> <li>Alcon Advisor and Speaker</li> <li>Amicus Advisor</li> <li>Aurelio Advisor</li> <li>Aura Advisor</li> <li>Bio Tissue Advisor and Speaker</li> <li>Bruder Advisor</li> <li>B&amp;L Advisor and Speaker</li> <li>Compe Advisory and Speaker</li> <li>Johnson and Johnson Advisor Speaker</li> <li>Kala Advisor and Speaker</li> <li>Lumena Advisor and Speaker</li> <li>Novartis Advisor and Speaker</li> <li>OutSoft Advisor</li> </ul>	<ul style="list-style-type: none"> <li>Occuphire Advisor</li> <li>Oyster Point Advisor and Speaker</li> <li>Orasol Advisor</li> <li>Ophthalmic Resource Partner</li> <li>Quidel Advisor</li> <li>RVA Advisor and Speaker</li> <li>Science Based Health Advisor and Speaker</li> <li>Sight Science Advisor and Speaker</li> <li>Sun Advisor and Speaker</li> <li>Tanus Advisor</li> <li>Thera Advisor</li> <li>Trafica Advisor</li> <li>Versa Advisor</li> <li>Vista Advisor/Global Advisor</li> </ul>
---	---

2




**Learning Objectives**

*Upon completion of this activity, the participant should be able to:*

- Summarize the etiologies of neurotrophic keratitis (NK) and how to differentiate it from similar diseases
- Recognize the newly proposed stages of NK
- Describe the stepwise approach to therapy and determine when to treat/refer patients
- Review clinical data on new and emerging treatments for NK


3

**Neurotrophic Keratitis and Staging**



4

**Neurotrophic Keratitis**




- NK is Classified as a Rare Disease
- Rare/orphan disease (ORPHA137596)<sup>1</sup>
- Affects < 5 individuals in 10,000
- NK Prevalence difficult to determine<sup>1,2</sup>
- Estimated to be < 1.6/10,000
- Best data are based on extrapolation from the most common conditions associated with NK
  - Herpes simplex keratitis: 6% develop NK
  - Herpes zoster keratitis: 12.8% develop NK
  - Postsurgical nerve damage: 2.8% develop NK

1. Devries DK, et al. Prog Retinal Eye Res. 2018;64:107-133.  
2. Sankar S, et al. Ocul Surf. 2011;13(4):299-310.

5

**Differential Diagnosis**


- Loss of corneal sensation = NK
- Neuropathic pain (corneal neuralgia, keratoneuralgia):
  - Pain without signs
  - Pain in response to minimal or even no stimulus
- Diseases with overlapping features of NK; can lead to NK if corneal sensation is affected<sup>1,2</sup>
  - Dry eye disease
  - Contact lens-related disorders
  - Blepharitis
  - Exposure keratopathy
  - Stem cell deficiency
  - Topical drug toxicity
  - Mild chemical injury



1. Devries DK, et al. Prog Retinal Eye Res. 2018;64:107-133.  
2. Sankar S, et al. Ocul Surf. 2011;13(4):299-310.

6

### Neurotrophic Keratitis Definition




- Degenerative corneal disease
- Damage to the trigeminal nerve (cranial nerve V)
- Loss of corneal sensation
- Breakdown of the corneal epithelium
- Impaired corneal healing
- Persistent epithelial defect → corneal ulceration → stromal melting and perforation

Hallmark: decreased sensation, decreased or no pain

7


### Corneal Innervation



- The cornea is the most sensitive and densely innervated tissue in the human body<sup>1,2</sup>
- Corneal innervation is essential. Corneal epithelial cells act in a mutually supportive relationship with corneal nerves<sup>1-4</sup>
  - Corneal nerves: maintain corneal integrity
  - Protective functions: blinking and tearing
  - Trophic support: neuropeptides (eg. substance P) promote epithelial cell proliferation, migration, adhesion
  - Epithelial cells: neurotrophic factors (neuronal extension and survival)
- Corneal nerve damage = loss of corneal sensation, epithelial breakdown, poor healing<sup>1,2</sup>

8

### Etiology




- INFECTIOUS<sup>1,2</sup>**
  - Herpes (simplex, zoster)
  - Leprosy
- ATROGENIC<sup>1,2</sup>**
  - Trauma to ciliary nerves by laser treatment and surgery
  - Corneal incisions
  - LASIK
- SYSTEMIC DISEASE<sup>1,2</sup>**
  - Diabetes
  - Multiple sclerosis
  - Vitamin A deficiency
- CORNEAL DYSTROPHIES<sup>1,2</sup>**
  - Lattice
  - Granular
- TOXIC<sup>1,2</sup>**
  - Chemical burns
  - Carbon disulfide exposure
  - Hydrogen sulfide exposure
- TOPICAL MEDICATIONS<sup>1,2</sup>**
  - Anesthetics (abuse)
  - Timolol
  - Betaxolol
  - Sulfacetamide
  - Diclofenac sodium
  - Ketorolac
- MSC<sup>1</sup>**
  - CTL
  - Increasing age
  - Adie syndrome
  - Limbal stem cell failure (chronic)
- FIFTH-NERVE PALSY<sup>1,2</sup>**
  - Trigeminal neuralgia surgery
  - Neoplasia (acoustic neuroma)
  - Aneurysms
  - Facial trauma
  - Congenital
  - Riley-Day syndrome
  - Goldenhar-Gorlin syndrome
  - Möbius syndrome
  - Familial corneal hypesthesia

9


### Chronic Comorbidities May Worsen Prognosis of NK

Chronic comorbidities can also confound the diagnosis of NK, increasing the need for a thorough diagnostic work-up, including a confirmatory test.



10

### Nerve Malfunction: Central to NK




11


### Etiologies: Impairment of Trigeminal Innervation

- Herpetic Corneal Disease (HSV/VZV)
- Damage to CN V - h/o stroke, tumor, brain injury/surgery
- H/o **LASIK** or other **ocular surgery**
- Iatrogenic injury (h/o contact lenses)
- Chronic use of topical medications (e.g., PGA timolol, betaxolol)
- Some corneal dystrophies
- Limbal stem cell deficiency (long standing/diseased epithelium (chemical burns))
- Systemic Diseases: ie, **diabetes mellitus**, multiple sclerosis, Riley-Day syndrome
- Multiple ocular surgeries
- Ocular cicatricial pemphigoid


12




13



14




15



16



17



18



19



20

**Handheld Esthesiometer (Cochet-Bonnet)**

**Steps:**

- Extend the retractable nylon monofilament to full length of 6 cm
- Retract the filament incrementally in 0.5 cm steps until the patient can feel its contact
- Record the length (shorter length indicates decreased sensation)
- Compare the fellow cornea
- Repeat steps 1-4 in each quadrant (superior, temporal, inferior, nasal)

21

**Mackie Severity Classification**

Stage	Clinical Features
1	<ul style="list-style-type: none"> <li>✓ Punctate epitheliopathy (punctate corneal fluorescein/LG staining)</li> <li>✓ Decreased TBUT</li> <li>✓ Stromal haze</li> </ul>
2	<ul style="list-style-type: none"> <li>✓ Persistent epithelial defect with smooth rolled edges</li> <li>✓ Stromal opacity</li> </ul>
3	<ul style="list-style-type: none"> <li>✓ Stromal thinning/ulceration</li> <li>✓ Corneal perforation</li> </ul>

Mackie, M. (1992) Neurotrophic keratitis. MB Saunders. Quack-Hil, Sand DG, Meunier DM, et al. Neurotrophic keratopathy. Prog Retin Eye Res. 2018;68:107-131.

22

Stage 1
Rose bengal staining of the inferior palpebral conjunctiva
Decreased TBUT
Increased mucous viscosity
Punctate corneal epithelial fluorescein staining (resembles dry eye)

**Mackie Classification**  
*Classified NK into 3 stages*

Mackie M in: Fourtaker C, Ray PG, Major DE eds. Current Ocular Therapy. WB Saunders 1992.


23

Stage 2
Epithelial defect
<ul style="list-style-type: none"> <li>• Typically oval in shape</li> <li>• In central/inferior cornea</li> <li>• Surrounded by a rim of loose epithelium</li> <li>• Edges may become smooth and rolled</li> </ul>
Stromal swelling with folds in the Descemet membrane
Anterior chamber inflammatory reaction may be present

**Mackie Classification**  
*Classified NK into 3 stages*

Mackie M in: Fourtaker C, Ray PG, Major DE eds. Current Ocular Therapy. WB Saunders 1992.

24




Stage 3
• Corneal ulcer
• Stromal lysis/melting
• Perforation

**Mackie Classification**  
*Classified NK into 3 stages*

Mackie Classification © Jay Tu, MD, Elmer Tu, MD, Linné Gustaf Therapeutics, MD, PhD, LLC


25



**Mackie Classification Summary**

- Commonly used in clinic and research
- Clustered a number of distinct and often nonsequential phases of NK development into 3 categories
- Very broad and nonspecific
- Recent advent of more effective treatment options necessitates a more highly defined staging system that better reflects the evolution of the disease and alerts clinicians to the earlier stages of NK

26



**The Neurotrophic Keratitis Study Group**

27

**The Neurotrophic Keratitis Study Group**

MEMBERS

- Edward J. Holland, MD - Chair
- Kenneth A. Beckman, MD
- Albert Y. Cheung, MD
- Marjan Farid, MD
- Nicole Fram, MD
- Preeya K. Gupta, MD
- W. Barry Lee, MD
- Francis S. Mah, MD
- Mark J. Mannis, MD
- Jay PePOSE, MD
- Elmer Tu, MD

- Proposed a new 7-step clinical staging system to more precisely classify the signs and symptoms associated with NK
- This classification will:
  - allow for earlier diagnosis
  - accurately monitor progression, evolution or recurrence
  - assess and evaluate its response to treatment

28


**Neurotrophic Keratitis Study Group Proposed Staging System**

Altered Sensation Without Keratopathy

- Patient can have absent sensation and not corneal findings

Stage 0 (Mild)


29




Stage 1 (Mild)

*Epitheliopathy without stromal haze*


30




31




32



33



34



35


### Topics for Discussion

- Mackie Classification versus Neurotrophic Keratitis Study Group 7-stage grading
  - When to use each?
- How to differentiate early NK from dry eye
  - Exam flow, staff flow
- When to manage?
- When to treat/refer?

36


**Diagnosis Conclusion:**  
Think NK

- History taking is a key component of patient assessment
- Rule out causes of impairment of trigeminal innervation
- Complete eye examination (epithelial defect may not be present! NK Mackie Stage 1)
- Corneal sensitivity testing
- Ancillary testing (Schirmer test, corneal cultures, confocal microscopy, r/o immune disorders)



37

**NK Treatment**



38

**Severity-Based Therapy**

Stage	Therapy
1	<ul style="list-style-type: none"> <li>• Preservative-free artificial tears formulations</li> <li>• Punctal occlusion</li> <li>• Hydrogel contact lens (consider large diameter)</li> <li>• Recombinant human NGF (rhNGF, cenegelein)</li> <li>• Serum/plasma/platelet rich plasma</li> </ul>
2	Supportive therapies plus: <ul style="list-style-type: none"> <li>• rhNGF</li> <li>• Scleral lens (± serum/plasma)</li> <li>• Amniotic membrane</li> <li>• Botulinum induced ptosis, tarsorrhaphy</li> </ul>
3	<ul style="list-style-type: none"> <li>• rhNGF</li> <li>• Keratoplasty + scleral lens, tarsorrhaphy, neurotization</li> </ul>

Santhi M, Lamborn A. Diagnosis and management of neurotrophic keratitis. Clin Ophthalmol. 2014;8:571-579. Shaha H, Tigha S, Hashem O, Hayashida Y. Update on keratinogen eye drops: the treatment of neurotrophic keratitis. Clin Ophthalmol. 2015;12:1012-1080. Published On: 2015

39

*Therapeutic Bandage Contact Lens*

**PROS**

- Inexpensive
- Mechanical protection
- Surface hydration

**CONS**

- Risks
  - Infection
  - Hypopyon formation
  - Reactive iritis
- **Requires frequent follow-up**
- Use with caution!

Alfon VO, Maitavola V. Management of NK. Contact Lens Adv Exp 2008;24:343-6. Maitavola VM, Maitavola V. Contact Lens Adv Exp 2008;24:319-9.

40

**Serum/Plasma Therapy**

**Serum/plasma have reported efficacy as primary or adjunct therapy**

- Reported success of serum alone (20-50% concentration) ranges from 71 to 100% within 90 days (Guadilla et al. Arch Soc Esp Ophthalmol 2013; Jeng and Dupps Cornea 2009; Pflugfelder AJO 2006)
- Umbilical cord serum may be more effective and has higher concentrations of substance P and NGF than peripheral blood serum (Yoon KC et al. Ophthalmology 2007)
- Epithelial defect healed in 97.4% of stage 2-3 NK after 11 weeks of plasma rich in growth factors (PRGF) (Sanchez-Avila RM et al. Int Ophthalmol 2018)
- Serum can be used safely in combination with SIH CL. No inflammation or CL deposits were observed (Choi JA ECL 2011)

41

**Amniotic Membrane**

- Randomized clinical trial reported healing of refractory neurotrophic ulcers with conventional therapy (lubrication plus BCL or tarsorrhaphy) or amniotic membrane transplant (AMT). Healing rates were similar in the 2 groups: 67% with conventional therapy and 73% with AMT (Khokhar S et al. Cornea 2005)
- AMT was also equivalent to autologous serum (AS) in healing neurotrophic ulcers: 70% for AS and 73% for AMT (Turkoglu E et al. Semin Ophthalmol 2014)
- Multilayer AMT recommended for deep ulcers and Descemetocelles (Kruse F et al. Ophthalmology 1999)

42

### Amniotic Membrane

- Self-retaining or in O.R.
- Single or multi-layer graft or patch
- Heal acute defect
- Restore stromal thickness
- Re-establish epithelial integrity
- Consider amniotic membrane extract




Image courtesy of Allergan Inc., MO.

43

### Lasting Effect by Increasing Corneal Nerve Density




Table 1. et al. Journal of Ophthalmology 2017 Aug 15; 2017.


44

### Scleral Lenses

- Use of fluid filled scleral contact lenses for treatment of NK initially reported decades ago (Romero-Rangel et al. AJO 2000)
- Nonhealing corneal epithelial defects with BCL healed without recurrence in all 9 eyes treated with PROSE scleral lens (Ling J et al. Am J Ophthalmol 2013)
- Overnight wear (with close monitoring) may accelerate healing (Lim P et al. AJO 2013)


45

### The Latest Treatment Options



47

### Active Ingredient Structurally Identical to Human Nerve Growth Factor Produced in Ocular Tissues




- Naturally occurring neurotrophin is responsible for differentiation, growth, and maintenance of neurons<sup>1</sup>
- The regenerative potential of nerve growth factor (NGF) was discovered by Nobel-prize winning scientists in the early 1950s<sup>1</sup>
- Cenegegermin-bkbj, a novel recombinant human nerve growth factor (rhNGF), is STRUCTURALLY IDENTICAL to the NGF protein<sup>2</sup>

<sup>1</sup> Linnemann R, Bergsøe P, Bostov S, Caporoglio G, Alami I. Topical treatment with nerve growth factor for corneal neurotrophic ulcers. N Engl J Med 1998;338:1174-80. <sup>2</sup> Vrabec R. New Drug Trends Report: Identifying Neurotrophic Treatments. AAOHN. 01/05/2010; 1(2):100.

48

### Endogenous NGF Maintains Corneal Integrity By Three Mechanisms

Endogenous Nerve growth factor acts through specific high-affinity (ie, TrkA) and low-affinity (ie, p75NTR) nerve growth factor receptors in the anterior segment of the eye to support corneal innervation and integrity.<sup>1</sup>



**SHOWN IN PRECLINICAL MODELS<sup>1</sup>**

**TEAR SECRETION**

**CORNEAL INNERVATION**

**CELL PROLIFERATION AND DIFFERENTIATION**


<sup>1</sup> Mithranathan S, Mithranathan S, Chakrabarti S, Nishida M, Sachdev M. Understanding the pathogenesis of neurotrophic keratitis: the role of corneal nerves. J Cell Physiol 2017;129:1712-24. <sup>2</sup> Miller L, Moller D, Kwan S, Tervo T. Corneal nerve structure, function, and function. Exp Eye Res 2003;76:252-61. <sup>3</sup> Sachdev M, Linnemann R. Changes and management of neurotrophic keratitis. Clin Ophthalmol 2014;8:217-8. <sup>4</sup> Alami I, Cullen-Weaver S, Corneal E, et al. Nerve Growth Factor in the Developing and Adult Corneal. Cornea of Regeneration and Repair.

49







50




51



52



53



54



55

### Recent Research

- Bruscolini et al performed a retrospective chart evaluation of 18 NK pts with at least 2 years f/u, n=10 at 36 mo, up to 48 mo (n=9).
- All 18 cleared at 8 weeks. At 1 year, 3 recurred. At 24 mo, 0 recurred. At 36 mo, 1/10 recurred. At 48 mo 0/9 recurred. VA, corneal sensitivity and tear production showed statistically significant differences at 1, 2 and 3 years (*Journal of Rare Diseases 2022*).

56

### Recent Research

- Pedrotti et al performed a prospective case series, n=18, 14/18 cleared at 8 weeks and stayed clear at 4 and 8 mo follow-up. In vivo corneal microscopy was used to evaluate corneal nerve regeneration.
- Significant peripheral corneal nerve growth and branching was seen at 2 mo, and central advancement across the 8 months. Corneal sensitivity improved. The nerve regeneration was partially visible at 8 weeks and continued after treatment with the hypothesis that the initial growth sustained further regeneration (*Journal of Rare Diseases 2022*).

57

### Recent Research

- Bonzano et al evaluated anterior segment OCT in 16 NK patients, half treated with 50% autologous serum and half with cenegermin.
- The corneal wound healing process was followed, including size and depth measured at the thinnest part of the cornea. Mean time to wound closure (slit lamp) was 3.9 weeks +/- 0.5 weeks and 5.9 weeks +/- 1.9 weeks in the AS arm.
- AS-OCT healing process: corneal epithelial hypertrophy, opaque reflective scar tissue followed by improvements in stromal thickness.
- Both treatments both improved NK, but cenegermin resolved quicker, possibly due to peripheral nerve regeneration. (*Frontiers in Pharmacology 2022*)

58

### Recent Research

- Sacchetti et al evaluated 2 groups, Amniotic membrane transplant and cenegermin with 12 months f/u: 13/15 AMT and 23/24 cenegermin remained cleared. There was less recurrence in the cenegermin group.
- Patient satisfaction and satisfaction with treatment outcomes were significantly better in the cenegermin group using a specifically designed patient reported satisfaction questionnaire.
- Similar to other studies, there was approximately a 13% recurrence rate. Survival analysis (recurrence) favored cenegermin. BCVA was statistically significantly improved.

59

### Treatment Summary

- Neurotrophic keratopathy is caused by a number of conditions
- Severity ranges from diffuse epitheliopathy to corneal ulceration and perforation
- Base treatment on severity stage
- Efficacy of many therapies are based on low level of evidence
- rhNGF is a validated, highly effective FDA-approved therapy that should be considered a first-line option
- A proactive approach to minimize recurrent corneal epithelial breakdown, stromal scarring and thinning and vision loss is recommended

61

## CASE STUDIES

62

**Case 1: LASIK NK Case**


Patient Information	<ul style="list-style-type: none"> <li>53-year-old woman</li> <li>Works in our billing office and sits in front of a computer all day</li> </ul>
Medical History	<ul style="list-style-type: none"> <li>Hx: LASIK Q4.2017</li> <li>Hx right side trigeminal neuralgia, 6-2017 had rhizotomy which did not help but resulted in right side facial and eye numbness</li> <li>Complains of decreased vision                             <ul style="list-style-type: none"> <li>As the day progresses, her central more than peripheral vision becomes hazy</li> <li>Uses artificial tears and notices it helps her vision for a brief period</li> <li>Feels no pain</li> <li>Later in the day, the vision is so bad she just covers the right eye</li> </ul> </li> <li>Being referred for a large central corneal abrasion OD</li> </ul>

Case courtesy of Francis Moh, MD

63

**Case 1: LASIK NK Case**


Rx	<ul style="list-style-type: none"> <li>Currently using ciprofloxacin 3-4x/day</li> </ul>
VA	<ul style="list-style-type: none"> <li>OD: 20/40 ph no improvement</li> <li>OS: 20/40 ph 20/20</li> </ul>



64

**Case 1: LASIK NK Case**


- Healed within 2 weeks using ointment QID OD



65

**Case 1: LASIK NK Case**

- However, during the next 12 months, every time she stopped the ointment, she would form another abrasion. She didn't like the ointment because it blurred her vision.
- She developed an abrasion 4 times within the year.
- Self-retaining AMT was used; ointment was used, but she kept breaking down when she decreased the ointment use.
- She was fitted for a scleral lens, but she couldn't tolerate it.
- Finally, we discussed tarsorrhaphy.



66


**Case 1: LASIK NK Summary**

- cenegermin launched in early 2019
- 1/28/2019 we prescribed cenegermin 6 x a day OD
- 2/11/2019 she was approved by her insurance
- 2/20/2019 she started cenegermin
- 2/21/2019 she saw the oculo-plastic surgeon to have the tarsorrhaphy taken down
- 3/20/2019 she was already healed
- 6/3/2022 she remains healed on artificial tears; VA 20/25

67

**Case 2**




Patient Information	<ul style="list-style-type: none"> <li>75-year-old man with 3- to 4-month nonhealing epithelial defect</li> </ul>
Medical History	<ul style="list-style-type: none"> <li>h/o bilateral LASIK</li> <li>h/o Herpes Zoster Ophthalmicus</li> <li>1 previous history of "corneal abrasion" 1 year ago that healed after 2 weeks with aggressive lubrication, antibiotic gts</li> </ul>
Previous Treatments	<ul style="list-style-type: none"> <li>BCL</li> <li>Amniotic Membrane (self retaining) - Prokera x 2</li> <li>Autologous serum gts</li> </ul>
Concomitant Medications	<ul style="list-style-type: none"> <li>Antibiotic gts</li> <li>Artificial tears</li> <li>Valtrex 1 gm BID</li> </ul>
Corneal Sensitivity	<ul style="list-style-type: none"> <li>Absent</li> </ul>
Diagnosis	<ul style="list-style-type: none"> <li>Nonhealing neurotrophic corneal epithelial defect</li> </ul>



Case courtesy of Meyer-Rand, MD

68


### Case 2—what to expect

Baseline	Week 4	Week 8
<ul style="list-style-type: none"> <li>&gt;5 mm central lesion, started cenegermin-bbj</li> </ul> 	<ul style="list-style-type: none"> <li>Central lesion reduced in size, incomplete closure</li> </ul> 	<ul style="list-style-type: none"> <li>Central lesion resolved, slight haze</li> </ul> 

cenegermin-bbj clinical trial

69




### Case 3

Patient Information	<ul style="list-style-type: none"> <li>84-year-old woman referred for ocular surface evaluation</li> </ul>	 <p>Case courtesy of Walter G. Whitely, OD, MBA, FFAO</p>
Medical History	<ul style="list-style-type: none"> <li>Dry eye syndrome - 10 years</li> <li>Hx herpes stromal keratitis</li> <li>Hx anterior scleritis</li> <li>POAG - mild s/p iStent OU</li> <li>NDDM</li> <li>Hypothyroid</li> <li>Seasonal allergies</li> </ul>	
Previous Treatments	<ul style="list-style-type: none"> <li>Punctal cautery</li> <li>Cryopreserved Amniotic Membrane (self retaining) x 2 OS</li> </ul>	
Concomitant Medications	<ul style="list-style-type: none"> <li>Aggressive preservative free artificial tears</li> <li>Cyclosporine 0.05%</li> </ul>	
Corneal Sensibility	<ul style="list-style-type: none"> <li>Central cornea decreased sensation, absent centrally</li> </ul>	
Diagnosis	<ul style="list-style-type: none"> <li>Nonhealing punctate keratopathy OS&gt;OD Stage 1 NK (severe)</li> </ul>	

70

### Case 3

Treated with cenegermin for 8 weeks

Baseline	Week 4	Week 8
		

- The most common adverse event (AE) seen in clinical trials was eye pain in approximately 16% of patients
- Other AEs occurring in 1% to 10% of patients and more frequently than in the vehicle-treated patients included corneal deposits, foreign body sensation, ocular hyperemia, ocular inflammation, and tearing

Observations are based on actual patient 112 from a cenegermin-bbj clinical trial.

71

THANK YOU!

73